

Appendix A.

The Methamphetamine Treatment Project

Overview

Conducted over 18 months between 1999 and 2001, the Methamphetamine Treatment Project (MTP) is (to date) the largest randomized clinical trial of treatment approaches for methamphetamine dependence; 978 individuals participated in the study (Rawson et al. 2004). MTP researchers randomly assigned participants at each treatment site into either the Matrix model treatment or the program's treatment as usual (TAU). The study design did not standardize TAU across sites, so each program offered different outpatient treatment models (including lengths of treatment ranging from 4 to 16 weeks). All TAU

models, along with the Matrix model, either required or recommended that participants attend 12-Step or mutual-help groups during their treatment, and all treatment models encouraged participation in continuing care activities after primary treatment.

The characteristics of a cross-section of participants in MTP (both TAU and Matrix participants) were found to be consistent with those of the clinical populations who participated in similar studies of treatment for methamphetamine abuse (Huber et al. 1997; Rawson et al. 2000). Figure A-1 lists specific client characteristics.

Figure A-1. Characteristics of MTP Participants			
Male	45%	Average education	12.2 years
Female	55%		
Caucasian	60%	Employed	69%
Hispanic/Latino	18%		
Asian/Pacific Islander	17%		
Other*	5%		
Average age	32.8 years	Average lifetime methamphetamine use	7.54 years
		Average days of methamphetamine use in the past 30 days	11.53 days
Married and not separated	16%	Preferred route of methamphetamine administration	
		Smoking	65%
		Intravenous	24%
		Intranasal	11%

*Two percent of participants in the Other category were African American (personal correspondence with Jeanne Obert, Matrix Institute, November 2004).

Source: Rawson et al. 2004, p. 711.

Participants' histories indicated multiple substance use. During the study, participant self-reports and drug and breath-alcohol tests confirmed that some clients had used marijuana or alcohol, as well as methamphetamines, but virtually no other substances of abuse were identified.

All MTP participants completed baseline assessments including the methamphetamine-dependence checklist in the *Diagnostic and Statistical Manual of Mental Disorders*, Fourth Edition (American Psychiatric Association 1994), and the Addiction Severity Index (McLellan et al. 1992). The assessments were repeated at several points during participants' active treatment, at discharge from treatment, and at 6 and 12 months after their dates of discharge from the program. Urine drug testing was conducted weekly throughout active treatment.

Results

No significant differences in substance use and functioning were found between TAU and Matrix groups at discharge and at 6-month followup. However, the MTP study found that the Matrix model participants (Rawson et al. 2004)

- Had consistently better treatment retention rates than did TAU participants
- Were 27 percent more likely than TAU participants to complete treatment
- Were 31 percent more likely than TAU participants to have methamphetamine-free urine test results while in treatment

At 6-month followup, more than 65 percent of both Matrix and TAU participants had negative urine tests for methamphetamine and other drugs (Rawson et al. 2004).

Appendix B.

Notes on Group Facilitation

All clients in a group, even in the highly structured Family Education group, develop individual relationships with their counselor. The degree to which the counselor can instigate positive change in clients' lives and support their family members is related directly to the credibility that he or she establishes. The counselor must be perceived as a highly credible source of information about substance use. Two keys to establishing credibility with clients are the degree to which the counselor engages and maintains control over a group and the counselor's ability to make all participants perceive the group as a safe place.

These two elements are highly interrelated. For a group to feel safe, the members need to view the counselor as competent and in control. Sometimes, group members enter the group with a lot of energy and are talkative and boisterous. Frequently this situation occurs during holidays, particularly if several members have relapsed. The counselor should use verbal and nonverbal methods of calming the group and focusing the group on the session topic. Conversely, there may be times when group members are lethargic, sluggish, and depressed. During these times, the counselor should infuse energy and enthusiasm. He or she needs to be aware of the emotional tone of families and respond accordingly, particularly when working with multifamily groups.

Because of the effects substance use disorders have on family members, negative emotions may be very close to the surface. The counselor must be prepared to handle these emotions appropriately as they arise and not to allow negative discussions or arguments to dominate a group session.

The members of a group need to feel that the counselor is keeping the group moving in a useful and healthful direction. The counselor must be willing to interrupt private conversations in the group, terminate a graphic drug use story, or redirect a lengthy tangential diversion. He or she must be perceived as clearly in control of the time in the group. Each member must be given an opportunity to have input. The counselor should ensure that a few members do not monopolize the group's time.

The counselor needs to be sensitive to emotional and practical issues that arise in group. At times it also may be necessary to be directive and confrontational or to characterize input from group members as a reflection of addictive thinking. In these instances, the counselor should focus on the addiction as opposed to the person. In other words, care should be taken to avoid directing negative feedback toward the client, focusing instead on the addiction-based aspects of the client's behavior or thinking.

The Family Education group counselor may be the professional who sees the clients for prescribed Individual/Conjoint sessions. The advantage of this dual role (group leader and individual counselor) is that the counselor can coordinate more effectively and guide the progressive recovery of each individual. The frequency of contact also strengthens the therapeutic bond that can hold the client in treatment. A potential disadvantage of the dual role is the possible danger that the counselor may inadvertently expose confidential client information to the group before the client chooses to do so. It is a violation of boundaries for the counselor even to imply that

information exists and to attempt to coerce a client into sharing that information if the client has not planned to do so in the group.

Another danger to be avoided is the counselor's being perceived as showing preference to some clients. It is important that the counselor be equally supportive of all group members and not allow them to engage in competition for attention.

The counselor can find discussions of group development, leadership, concepts, techniques, and other helpful information for conducting group therapy in Treatment Improvement Protocol 41, *Substance Abuse Treatment: Group Therapy* (CSAT 2005), a free publication from the Center for Substance Abuse Treatment.

Appendix C.

Acronyms and Abbreviations List

AA	Alcoholics Anonymous
ACoA	Adult Children of Alcoholics
Al-Anon	A support group for families and loved ones of people who are addicted to alcohol
Alateen	A support group for young family members and loved ones of people who are addicted to alcohol
ASI	Addiction Severity Index
CA	Cocaine Anonymous
CAL	Calendar (for worksheets used during scheduling)
CMA	Crystal Meth Anonymous
CoDA	Co-Dependents Anonymous
CSAT	Center for Substance Abuse Treatment
EA	Emotions Anonymous
ERS	Early Recovery Skills
GA	Gamblers Anonymous
HALT	Hungry Angry Lonely Tired
IC	Individual/Conjoint
IOP	Intensive Outpatient Treatment for People With Stimulant Use Disorders
JACS	Jewish Alcoholics, Chemically Dependent Persons and Significant Others
MA	Marijuana Anonymous
meth	Methamphetamine
MTP	Methamphetamine Treatment Project
NA	Narcotics Anonymous
Nar-Anon	A support group for families and loved ones of people who are addicted to narcotics
OA	Overeaters Anonymous
PA	Pills Anonymous
RP	Relapse Prevention
SAMHSA	Substance Abuse and Mental Health Services Administration
SCH	Schedule (for worksheets used during scheduling)
SMART	Self-Management and Recovery Training
SS	Social Support
TAU	Treatment as Usual

Appendix D.

Field Reviewers

Rosie Anderson-Harper, M.A., RSAP
Mental Health Manager
Missouri Department of Mental Health
Jefferson City, MO

Stephen R. Andrew, M.S.W., LCSW,
LADC, CGP
Director
Health Education Training Institute
Portland, ME

Michelle M. Bartley
Behavioral Health Specialist
Division of Behavioral Health
Anchorage, AK

Frances Clark, Ph.D., MAC, LADAC,
QSAP, CCJS
Director of Behavioral Services
Metro Public Health Department
Nashville, TN

María del Mar García, M.H.S., LCSW
Continuing Education Coordinator
Caribbean Basin and Hispanic Addiction
Technology Transfer Center
Universidad Central del Caribe
Bayamón, PR

Darcy Edwards, Ph.D., M.S.W., CADC II
Substance Abuse Treatment Coordinator
Oregon Department of Corrections
Salem, OR

Marty Estrada, CAS, CSS-III
Ventura, CA

Eric Haram, LADAC
Administrative Specialist
Mercy Recovery Center
Westbrook, ME

Sherry Kimbrough, M.S., NCAC
Vice President
Lanstat, Inc.
Port Townsend, WA

Thomas A. Peltz, LMHC, LADAC-1
Therapist
Private Practice
Beverly Farms, MA

John L. Roberts, M.Ed., CCDC III-E, LPC, MAC
Consultant/Trainer
Continuing Education Center
Cincinnati, OH

Jim Rowan, M.A., LAC
Program Manager
Arapahoe House, Inc.
Thornton, CO

Angel Velez, CASAC
Addiction Program Specialist II
Office of Alcohol and Substance Abuse
Services
New York, NY

Appendix E.

Bibliography

- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*, Fourth Edition. Washington, DC: American Psychiatric Association, 1994.
- Center for Substance Abuse Prevention. *Ecstasy: What's All the Rave About?* DHHS Publication No. (SMA) 3710B. Rockville, MD: Substance Abuse and Mental Health Services Administration, 2002.
- Center for Substance Abuse Prevention. *Marijuana: Weeding Out the Hype*. DHHS Publication No. (SMA) 3710D. Rockville, MD: Substance Abuse and Mental Health Services Administration, 2002.
- Center for Substance Abuse Prevention. *Heroin: What's the Real Dope?* DHHS Publication No. (SMA) 3710C. Rockville, MD: Substance Abuse and Mental Health Services Administration, 2003.
- CSAT (Center for Substance Abuse Treatment). *Treatment for Stimulant Use Disorders*. Treatment Improvement Protocol (TIP) Series 33. DHHS Publication No. (SMA) 99-3296. Rockville, MD: Substance Abuse and Mental Health Services Administration, 1999.
- CSAT (Center for Substance Abuse Treatment). *Substance Abuse Treatment: Group Therapy*. Treatment Improvement Protocol (TIP) Series 41. DHHS Publication No. (SMA) 05-3991. Rockville, MD: Substance Abuse and Mental Health Services Administration, 2005.
- CSAT (Center for Substance Abuse Treatment). *Improving Cultural Competence in Substance Abuse Treatment*. Treatment Improvement Protocol (TIP) Series. Rockville, MD: Substance Abuse and Mental Health Services Administration, forthcoming.
- Graham, A.W.; Schultz, T.K.; Mayo-Smith, M.F.; Ries, R.K.; and Wilford, B.B., eds., *Principles of Addiction Medicine*, Third Edition. Chevy Chase, MD: American Society of Addiction Medicine, Inc., 2003.
- Huber, A.; Ling, W.; Shoptaw, S.; Gulati, V.; Brethen, P.; and Rawson, R. Integrating treatments for methamphetamine abuse: A psychosocial perspective. *Journal of Addictive Diseases* 16(4):41–50, 1997.
- Johnston, L.D.; O'Malley, P.M.; Bachman, J.G.; and Schulenberg, J.E. *Monitoring the Future National Results on Adolescent Drug Abuse: Overview of Key Findings 2004*. NIH Publication No. 05-5726. Bethesda, MD: National Institute on Drug Abuse, 2005. www.monitoringthefuture.org/pubs/monographs/overview2004.pdf [accessed September 16, 2005].

- Khalsa, J.H.; Genser, S.; Francis, H.; and Martin, B. Clinical consequences of marijuana. *Journal of Clinical Pharmacology* 42 (Suppl):7S–10S, 2002.
- McLellan, A.T.; Kushner, H.; Metzger, D.; Peters, R.; Smith, L.; Grissom, G.; Pettinati, H.; and Argeriou, M. The fifth edition of the Addiction Severity Index. *Journal of Substance Abuse Treatment* 9:199–213, 1992.
- National Institute on Alcohol Abuse and Alcoholism. Are women more vulnerable to alcohol's effects? *Alcohol Alert* 46, December 1999 (updated October 2000). www.niaaa.nih.gov/publications/aa46.htm [accessed September 16, 2005].
- National Institute on Alcohol Abuse and Alcoholism. Fetal alcohol exposure and the brain. *Alcohol Alert* 50, December 2000 (updated April 2001). www.niaaa.nih.gov/publications/aa50.htm [accessed September 16, 2005].
- National Institute on Alcohol Abuse and Alcoholism. *Helping Patients Who Drink Too Much: A Clinician's Guide*, 2005 Edition. Bethesda, MD: National Institutes of Health, 2005. www.niaaa.nih.gov/publications/Practitioner/guide.pdf [accessed September 16, 2005].
- National Institute on Drug Abuse. *NIDA Community Drug Alert Bulletin: Methamphetamine*. Bethesda, MD: National Institutes of Health, 1998. www.drugabuse.gov/MethAlert/MethAlert.html [accessed September 16, 2005].
- National Institute on Drug Abuse. *Research Report Series: Marijuana Abuse*. NIH Publication No. 02-3859. Bethesda, MD: National Institutes of Health, October 2002. www.drugabuse.gov/PDF/RRMarijuana.pdf [accessed September 16, 2005].
- National Institute on Drug Abuse. *NIDA InfoFacts: Marijuana*. Bethesda, MD: National Institutes of Health, March 2004. www.drugabuse.gov/PDF/InfoFacts/Marijuana04.pdf [accessed September 16, 2005].
- National Institute on Drug Abuse. *Research Report Series: Cocaine Abuse and Addiction*. NIH Publication No. 99-4342. Bethesda, MD: National Institutes of Health, revised November 2004. www.nida.nih.gov/PDF/RRCocain.pdf [accessed September 16, 2005].
- National Institute on Drug Abuse. *NIDA InfoFacts: LSD*. Bethesda, MD: National Institutes of Health, March 2005. www.nida.nih.gov/pdf/infofacts/LSD05.pdf [accessed September 16, 2005].
- National Institute on Drug Abuse. *NIDA InfoFacts: Rohypnol and GHB*. Bethesda, MD: National Institutes of Health, March 2005. www.drugabuse.gov/PDF/Infofacts/Rohypnol05.pdf [accessed September 16, 2005].
- Obert, J.; McCann, M.J.; Marinelli-Casey, P.; Weiner, A.; Minsky, S.; Brethen, P.; and Rawson, R. The Matrix model of outpatient stimulant abuse treatment: History and description. *Journal of Psychiatric Drugs* 32(2):157–164, 2000.

- Office of National Drug Control Policy. *Fact Sheet: Gamma Hydroxybutyrate (GHB)*. Rockville, MD: Drug Policy Information Clearinghouse, November 2002. www.whitehousedrugpolicy.gov/pdf/gamma_hb.pdf [accessed September 16, 2005].
- Office of National Drug Control Policy. *Pulse Check: Trends in Drug Abuse*. Washington, DC: Executive Office of the President, November 2002.
- Office of National Drug Control Policy. *Fact Sheet: Heroin*. Rockville, MD: Drug Policy Information Clearinghouse, June 2003. www.whitehousedrugpolicy.gov/publications/factsht/heroin/197335.pdf [accessed September 16, 2005].
- Office of National Drug Control Policy. *Fact Sheet: Methamphetamine*. Rockville, MD: Drug Policy Information Clearinghouse, November 2003. www.whitehousedrugpolicy.gov/publications/pdf/ncj197534.pdf. [accessed September 16, 2005].
- Office of National Drug Control Policy. *Fact Sheet: Rohypnol*. Rockville, MD: Drug Policy Information Clearinghouse, February 2003. www.whitehousedrugpolicy.gov/publications/factsht/rohypnol/rohypnol.pdf [accessed September 16, 2005].
- Office of National Drug Control Policy. *Fact Sheet: Marijuana*. Rockville, MD: Drug Policy Information Clearinghouse, February 2004. www.whitehousedrugpolicy.gov/publications/pdf/ncj198099.pdf [accessed September 16, 2005].
- Rawson, R.; Huber, A.; Brethen, P.; Obert, J.; Gulati, V.; Shoptaw, S.; and Ling, W. Methamphetamine and cocaine users: Difference in characteristics and treatment retention. *Journal of Psychoactive Drugs* 32(2):233–238, 2000.
- Rawson, R.A.; Marinelli-Casey, P.; Anglin, M.D.; Dickow, A.; Frazier, Y.; Gallagher, C.; Galloway, G.P.; Herrell, J.; Huber, A.; McCann, M.J.; Obert, J.; Pennell, S.; Reiber, C.; Vandersloot, D.; and Zweben, J. A multi-site comparison of psychosocial approaches for the treatment of methamphetamine dependence. *Addiction* 99(6):708–717, 2004.
- Rawson, R.A.; Shoptaw, S.J.; Obert, J.L.; McCann, M.J.; Hasson, A.L.; Marinelli-Casey, P.J.; Brethen, P.R.; and Ling, W. An intensive outpatient approach for cocaine abuse treatment: The Matrix model. *Journal of Substance Abuse Treatment* 12(2):117–127, 1995.
- Shoptaw, S.; Rawson, R.A.; McCann, M.J.; and Obert, J.L. The Matrix model of outpatient stimulant abuse treatment: Evidence of efficacy. *Journal of Addictive Diseases* 13(4):129–141, 1994.
- Substance Abuse and Mental Health Services Administration (SAMHSA). *Club Drugs, 2002 Update. The Dawn Report*. Rockville, MD: Office of Applied Studies, SAMHSA, July 2004.
- Substance Abuse and Mental Health Services Administration (SAMHSA). *The NSDUH Report: Daily Marijuana Users*. Rockville, MD: Office of Applied Studies, SAMHSA, November 26, 2004. www.oas.samhsa.gov/2k4/dailyMJ/dailyMJ.htm [accessed September 16, 2005].

Substance Abuse and Mental Health Services Administration (SAMHSA). *Overview of Findings From the 2003 National Survey on Drug Use and Health*. NSDUH Series H-24, DHHS Publication No. (SMA) 04-3963. Rockville, MD: Office of Applied Studies, SAMHSA, 2004.
oas.samhsa.gov/NHSDA/2k3NSDUH/2k3OverviewW.pdf [accessed September 16, 2005].

Substance Abuse and Mental Health Services Administration (SAMHSA). *Results From the 2003 National Survey on Drug Use and Health: National Findings*. NSDUH Series H-25, DHHS Publication No. (SMA) 04-3964. Rockville, MD: Office of Applied Studies, SAMHSA, 2004.

Substance Abuse and Mental Health Services Administration (SAMHSA). *The NSDUH Report: Methamphetamine Use, Abuse, and Dependence: 2002, 2003, and 2004*. Rockville, MD: Office of Applied Studies, SAMHSA, September 16, 2005. www.oas.samhsa.gov/2K5/meth/meth.pdf [accessed March 2, 2006].

Substance Abuse and Mental Health Services Administration (SAMHSA). *Results From the 2004 National Survey on Drug Use and Health: National Findings*. NSDUH Series H-28, DHHS Publication No. (SMA) 05-4062. Rockville, MD: Office of Applied Studies, SAMHSA, 2005.

Tevisan, L.; Boutros, N.; Petrakis, I.; and Krystal, J.H. Complications from alcohol withdrawal. *Alcohol Health and Research World* 22(1):61–66, 1998.



DHHS Publication No. (SMA) 06-4153
U.S. Department of Health and Human Services
Substance Abuse and Mental Health Services Administration